

This is an official

CDC Health Advisory

Distributed via Health Alert Network
March 18, 2009 15:00 EST (03:00PM EST)

Invasive *Haemophilus influenzae* Type B Disease in Young Children and Importance for All Young Children to Receive 3 Dose Primary Series with Available Hib-containing Vaccine

Background of the Hib Vaccine Shortage

Health care providers must be vigilant about ensuring that all young children are appropriately vaccinated with the 3 dose primary series of Hib (*Haemophilus influenzae* type b) vaccine. A nationwide shortage of Hib vaccine began in December 2007 and is ongoing. The recall of certain lots of the two Hib-containing vaccines produced by Merck & Co., Inc. and cessation of production of both vaccines has left only one manufacturer of Hib vaccine in the United States (sanofi pasteur). The shortage resulted in a recommendation by CDC to defer the Hib booster (routinely recommended at 12 through 15 months) for children who are NOT at high risk of Hib infection temporarily, until supplies are restored. This recommendation is still in effect.

Temporary deferral of the booster dose at 12 through 15 months of age for non-high risk children may have resulted in increased Hib carriage and transmission in non-symptomatic children. There is potential to see increases in cases of Hib disease at the local level. During 2008 in Minnesota, five children aged 5 months through 3 years were reported with invasive Hib disease; one died. Three patients had received no vaccinations because of parent or guardian deferral or refusal. One child was aged 5 months and had received 2 doses of Hib PRP-TT vaccine in accordance with the primary series schedule. Another child had received 2 doses of Hib PRP-OMP vaccine, but no booster dose, per CDC recommendations during the shortage. Subsequent to Hib infection, this child was diagnosed with hypogammaglobulinemia. The five cases in 2008 were the most reported for 1 year from Minnesota since 1992, when 10 cases were reported.

There is enough Hib-containing vaccine for all U.S. children to receive the primary series. All children should complete the primary series by 7 months of age; high risk children should continue to receive the full primary series and the booster dose. Completion of the primary series with currently available vaccine products (manufactured by sanofi pasteur) requires a total of 3 doses of Hib-containing vaccine (2, 4, and 6 months). Although there is enough Hib-containing vaccine nationally to support these recommendations, there may be times when practitioners do not have an adequate supply of vaccine to meet local demand. If Hib vaccine is not available in the office at the time of a visit, children who are unable to receive one of the primary series doses should be tracked and recalled to schedule an appointment to receive their dose as soon as vaccine becomes available in the office.

In addition, using available Hib-containing vaccines has presented challenges associated with switching from the Merck to sanofi products for some providers.

There are indications that these challenges have led to lower completion of the primary series. Preliminary information comes from sentinel immunization information systems (registries) in select states, which have indicated up to 10% lower coverage with the third Hib dose in the primary series compared to other vaccines (DTaP, PCV7) commonly administered at the same visit. In the scenario of booster dose deferral, it is even more important that all infants receive the complete primary series.

Specifically, some of the challenges in using the currently available Hib-containing vaccines have included provider reluctance to switch inventory and schedules, misunderstanding regarding what constitutes primary versus booster doses, determining a catch-up schedule in the setting of the deferred booster, and provider and

parent concerns about over vaccination resulting from switching to the sanofi pasteur Hib-containing vaccine. Despite these challenges, health care providers need to ensure that all children are appropriately vaccinated with the primary series. For example, if Pentacel (DTaP-IPV/Hib) is the only Hib-containing vaccine available, this combination product should be used to complete the primary series, even if doing so results in receipt of additional doses of other antigens (e.g., DTaP, IPV). The Hib-containing vaccine products that are available may not be what providers used previously in their practice; however, the potential for increased transmission of Hib makes it more important than ever that every child is adequately protected.

Recommendations

The following non-high risk children should be scheduled to receive the primary series of Hib vaccine as outlined below:

- If the child is at least 6 weeks but less than 12 months of age and has received zero, one, or two doses of Hib vaccine, schedule him/her for the first or next dose(s) immediately with a minimum of four weeks between the doses. These children will need one booster dose when the Hib vaccine shortage is over.
- If the child is between 12 and 14 months of age and has not had any doses of Hib vaccine, schedule appointments for two doses, eight weeks apart.
- If the child is between 12 and 14 months of age and has received Hib vaccine but did not complete the primary series before they turned 1 year old (i.e., had 1 dose of the Merck product OR 1-2 doses of the sanofi product), schedule an appointment for 1 additional dose, a minimum of eight weeks from the last dose.
- If the child is at least 15 months of age but less than 5 years of age and has not received any doses of Hib vaccine OR has not completed the primary series (i.e., had 1 dose of the Merck product OR 1-2 doses of the sanofi product), schedule an appointment for one dose.
- If the child is 5 years old or older and hasn't received any Hib vaccine, Hib vaccine is not necessary.

Certain children are at increased risk for Hib disease, including children with asplenia, sickle cell disease, human immunodeficiency virus infection and certain other immunodeficiency syndromes, and malignant neoplasms. CDC recommends that providers continue to vaccinate these children with available Hib conjugate vaccines according to the routinely recommended schedules, including the 12 through 15 month booster dose. Providers who serve predominantly American Indian/Alaska Native (AI/AN) children living in AI/AN communities should continue to stock and use PRP-OMP– containing Hib vaccines (Merck product) and vaccinate according to the routinely recommended schedule, which includes the 2-dose primary series (ages 2 and 4 months) and a 12 through 15 month booster dose. This product is available from the VFC Pediatric Vaccine Stockpile, through their state immunization programs.

Additional Sources of Information

For more information about Hib disease and vaccination contact your state or local public health official or CDC at 1-800-232-4636/1-800-CDC-INFO or by email at www.cdc.gov/vaccines/about/contact/nipinfo_contact_form.htm. Information about current vaccine shortages and delays can be found at <http://www.cdc.gov/vaccines/vac-gen/shortages/default.htm>.

- CDC. Invasive *Haemophilus influenzae* Type B Disease in Five Young Children – Minnesota, 2008. MMWR 2009;58:1-3.
- CDC. Continued shortage of *Haemophilus influenzae* Type B (Hib) Conjugate Vaccines and Potential Implications for Hib Surveillance – United States, 2008. MMWR 2008;57(46):1252-1255.
- CDC. Interim recommendations for the Use of *Haemophilus influenzae* Type B (Hib) Conjugate Vaccines Related to the Recall of Certain Lots of Hib-containing Vaccines (PedvaxHIB and Comvax). MMWR 2007;56(50):1318-1320.

DHEC Contact Information for Reportable Diseases and Reporting Requirements

Reporting of cases of *Haemophilus influenzae*, invasive disease (all types), is present on our current 2009 List of Reportable Conditions as Urgently Reportable by phone within 24 hours. Reporting is consistent with South Carolina Law requiring the reporting of diseases and conditions to your state or local public health department. (State Law # 44-29-10 and Regulation # 61-20) as per the DHEC 2009 List of Reportable Conditions available at: http://www.scdhec.gov/health/disease/docs/reportable_conditions.pdf

Federal HIPAA legislation allows disclosure of protected health information, without consent of the individual, to public health authorities to collect and receive such information for the purpose of preventing or controlling disease. (HIPAA 45 CFR §164.512).

Regional Public Health Offices – 2009

Mail or call reports to the Epidemiology Office in each Public Health Region.

Region 1

Anderson, Oconee

220 McGee Road
Anderson, SC 29625
Phone: (864) 260-4358
Fax: (864) 260-5623
Nights / Weekends: 1-866-298-4442

Abbeville, Edgefield, Greenwood, Laurens, McCormick, Saluda

1736 S. Main Street
Greenwood, SC 29646
Phone: 1-888-218-5475
Fax: (864) 942-3690
Nights / Weekends: 1-800-420-1915

Region 2

Greenville, Pickens

PO Box 2507
200 University Ridge
Greenville, SC 29602-2507
Phone: (864) 282-4139
Fax: (864) 282-4373
Nights / Weekends: 1-800-993-1186

Cherokee, Spartanburg, Union

PO Box 4217
151 E. Wood Street
Spartanburg, SC 29305-4217
Phone: (864) 596-2227, x- 210
Fax: (864) 596-3443
Nights / Weekends: 1-800-993-1186

Region 3

Chester, Lancaster, York

PO Box 817
1833 Pageland Highway
Lancaster, SC 29720
Phone: (803) 286-9948
Fax: (803) 286-5418
Nights / Weekends: 1-866-867-3886

Region 3 (continued)

Fairfield, Lexington, Newberry, Richland

2000 Hampton Street
Columbia, SC 29204
Phone: (803) 576-2749
Fax: (803) 576-2993
Nights / Weekends: 1-888-554-9915

Region 4

Clarendon, Kershaw, Lee, Sumter

PO Box 1628
105 North Magnolia Street
Sumter, SC 29150
Phone: (803) 773-5511
Fax: (803) 775-9941
Nights/Weekends: 1-877-831-4647

Chesterfield, Darlington, Dillon, Florence, Marlboro, Marion

145 E. Cheves Street
Florence, SC 29506
Phone: (843) 661-4830
Fax: (843) 661-4859
Nights / Weekends: (843) 660-8145

Region 5

Bamberg, Calhoun, Orangeburg

PO Box 1126
1550 Carolina Avenue
Orangeburg, SC 29116
Phone: (803) 533-7199
Fax: (803) 533-7134
Nights / Weekends: (803) 954-8513

Aiken, Allendale, Barnwell

1680 Richland Avenue, W. Suite 40
Aiken, SC 29801
Phone: (803) 642-1618
Fax: (803) 643-8386
Nights / Weekends: (803) 827-8668 or
1-800-614-1519

Region 6

Georgetown, Horry, Williamsburg

1931 Industrial Park Road
Conway, SC 29526-5482
Phone: (843) 915-8804
Fax: (843) 365-0085
Nights / Weekends: (843) 381-6710

Region 7

Berkeley, Charleston, Dorchester

4050 Bridge View Drive, Suite 600
N. Charleston, SC 29405
Phone: (843) 953-0060
Fax: (843) 953-0051
Nights / Weekends: (843) 219-8470

Region 8

Beaufort, Colleton, Hampton, Jasper

219 S. Lemacks Street
Walterboro, SC 29488
Phone: (843) 525-7603, x-108
Fax: (843) 549-6845
Nights / Weekends: 1-800-614-4698

DHEC Bureau of Disease Control Division of Acute Disease Epidemiology

1751 Calhoun Street
Box 101106
Columbia, SC 29211
Phone: (803) 898-0861
Fax: (803) 898-0897
Nights / Weekends: 1-888-847-0902



www.scdhec.gov

Categories of Health Alert messages:

Health Alert: conveys the highest level of importance; warrants immediate action or attention.

Health Advisory: provides important information for a specific incident or situation; may not require immediate action.

Health Update: provides updated information regarding an incident or situation; unlikely to require immediate action.